

On the Sturm–Liouville-Type Boundary Value Problem

B. G. PACHPATTE

*Department of Mathematics and Statistics, Marathwada University,
Aurangabad 431004 (Maharashtra), India*

Submitted by C. L. Dolph

The problems of existence, uniqueness and continuous dependence on parameter of solutions of the nonlinear boundary value problem of Sturm–Liouville type for ordinary differential equations are considered. The well-known Banach fixed point theorem is used to establish results. © 1985 Academic Press, Inc.

1. INTRODUCTION

Macki and Waltman [9] have studied the following Sturm–Liouville-type boundary value problem (BVP)

$$x' = f(t, x, y, \lambda), \quad (1)$$

$$y' = g(t, x, y, \lambda), \quad (2)$$

$$A_1 x(a) + A_2 y(a) = 0, \quad (3)$$

$$B_1 x(b) + B_2 y(b) = 0, \quad (4)$$

under some suitable conditions. The work of Macki and Waltman [9] examines the existence of eigenvalues of the parameter λ for which there exists a nontrivial solution of BVP (1)–(4). The problems of this type are of great importance from both the theoretical and the applied point of view (see [1, 6, 15]). Our interest in such problems was motivated by their occurrence in various branches of transport theory and in the problems of optimal economic growth; see [7, 8, 12, 13] and the references given therein. Here we assume throughout that $x, y \in C[I, E]$, $f, g \in C[I \times E \times E \times R, E]$, A_i, B_i ($i = 1, 2$) are constants such that $A_1 \neq 0$, $B_2 \neq 0$, where $I = [a, b]$, E denotes the Banach space with convenient norm $\|\cdot\|$ and R denotes the set of real numbers.

In this paper we present a set of conditions on functions f, g which are sufficient to guarantee that BVP (1)–(4) has a unique solution. We also

determine conditions for continuous dependence of solutions of BVP (1)–(4) on a parameter λ . Our results are obtained by using the Banach fixed point theorem. The sufficient conditions obtained here are different than those obtained by various authors in [2–5, 7–9, 12–14] and our approach to the problem is different as well.

2. MAIN RESULTS

In this section we establish our main results on the existence of a unique solution and continuous dependence on parameter of solutions of BVP (1)–(4). For convenience we first list the following hypotheses.

(H₁) For $\lambda \in R$, $(t, x_i, y_i, \lambda) \in I \times E \times E \times R$ ($i = 1, 2$),

$$\|f(t, x_1, y_1, \lambda) - f(t, x_2, y_2, \lambda)\| \leq p(t)[\|x_1 - x_2\| + \|y_1 - y_2\|],$$

$$\|g(t, x_1, y_1, \lambda) - g(t, x_2, y_2, \lambda)\| \leq q(t)[\|x_1 - x_2\| + \|y_1 - y_2\|],$$

where $p, q \in C[I, R^+]$ such that

$$\int_a^t p(s) \exp(Ls) ds \leq \alpha \exp(Lt),$$

$$\int_t^b q(s) \exp(Ls) ds \leq \beta \exp(Lt),$$

for $t \in I$, in which L is a positive constant and α, β are nonnegative constants such that $0 \leq \alpha + \beta < 1$.

(H₂) For every fixed $\lambda \in R$ there exist nonnegative constants N_1, N_2 such that

$$\int_a^t \|f(s, \theta, \theta, \lambda)\| ds \leq N_1 \exp(Lt),$$

$$\int_t^b \|g(s, \theta, \theta, \lambda)\| ds \leq N_2 \exp(Lt),$$

for $t \in I$, where θ is the zero element in E .

(H₃) There exist nonnegative constants M_1, M_2 and functions $p_0, q_0 \in C[I, R^+]$ such that for $(t, x, y, \lambda_i) \in I \times E \times E \times R$ ($i = 1, 2$),

$$\|f(t, x, y, \lambda_1) - f(t, x, y, \lambda_2)\| \leq p_0(t) |\lambda_1 - \lambda_2|,$$

$$\|g(t, x, y, \lambda_1) - g(t, x, y, \lambda_2)\| \leq q_0(t) |\lambda_1 - \lambda_2|,$$

and

$$\int_a^t p_0(s) ds \leq M_1 \exp(Lt),$$

$$\int_t^b q_0(s) ds \leq M_2 \exp(Lt),$$

for $t \in I$.

Let $B = E \times E$ be the product space equipped with the norm

$$\|z(t)\|_B = \|z_1(t)\| + \|z_2(t)\|, \quad z = (z_1, z_2) \in B, t \in I.$$

Let S be a space of those functions $z = (z_1, z_2) \in B$, which are continuous for $t \in I$ and fulfill the condition

$$\|z(t)\|_B = O(\exp(Lt)), \quad t \in I. \quad (5)$$

In the space S we define the norm (see [10])

$$\|z\|_1 = \sup_{t \in I} [\|z(t)\|_B \exp(-Lt)]. \quad (6)$$

It is easy to see that S with the norm defined in (6) is a Banach space.

We note that the condition (5) implies that there exists a constant $M \geq 0$ such that $\|z(t)\|_B \leq M \exp(Lt)$, $t \in I$. Using this fact in (6) we observe that

$$\|z\|_1 \leq M. \quad (7)$$

We now establish the following existence–uniqueness theorem for BVP (1)–(4).

THEOREM 1. *Assume that the hypotheses (H_1) and (H_2) hold. Then for every $\lambda \in R$ there exists a unique solution $(x, y) \in S$ of BVP (1)–(4).*

Proof. Let $\lambda \in R$ be fixed. For $z = (x, y) \in S$ we define the operator $Tz = (T_1 x, T_2 y)$, where the operators $T_1, T_2: E \rightarrow E$ are defined by

$$T_1 x(t) = c_1 + \int_a^t f(s, x(s), y(s), \lambda) ds, \quad (8)$$

$$T_2 y(t) = c_2 - \int_t^b g(s, x(s), y(s), \lambda) ds, \quad (9)$$

where $c_1 = -(A_2/A_1) y(a)$ and $c_2 = -(B_1/B_2) x(b)$. Clearly the solution of BVP (1)–(4) is a fixed point of the operator equation $Tz = z$. We first prove that T maps S into itself. Evidently T is continuous and $Tz \in E \times E$ for $t \in I$.

We verify that (5) is fulfilled. From (8) and (9) and using (H_1) and (H_2) and (7) we obtain

$$\begin{aligned}
 \|T_1 x(t)\| &\leq \|c_1\| + \int_a^t \|f(s, x(s), y(s), \lambda) - f(s, \theta, \theta, \lambda)\| ds \\
 &\quad + \int_a^t \|f(s, \theta, \theta, \lambda)\| ds \\
 &\leq \|c_1\| + \int_a^t p(s) \|z(s)\|_B \exp(-Ls) \exp(Ls) + N_1 \exp(Lt) \\
 &\leq \|c_1\| + \|z\|_1 \alpha \exp(Lt) + N_1 \exp(Lt) \\
 &\leq [\|c_1\| + M\alpha + N_1] \exp(Lt),
 \end{aligned} \tag{10}$$

and

$$\begin{aligned}
 \|T_2 y(t)\| &\leq \|c_2\| + \int_t^b \|g(s, x(s), y(s), \lambda) - g(s, \theta, \theta, \lambda)\| ds \\
 &\quad + \int_t^b \|g(s, \theta, \theta, \lambda)\| ds \\
 &\leq \|c_2\| + \int_t^b q(s) \|z(s)\|_B \exp(-Ls) \exp(Ls) ds + N_2 \exp(Lt) \\
 &\leq \|c_2\| + \|z\|_1 \beta \exp(Lt) + N_2 \exp(Lt) \\
 &\leq [\|c_2\| + M\beta + N_2] \exp(Lt).
 \end{aligned} \tag{11}$$

From (10) and (11) we have

$$\|Tz\|_1 \leq [\|c_1\| + \|c_2\| + M(\alpha + \beta) + N_1 + N_2].$$

This proves that T maps S into itself.

Now we verify that the operator T is a contraction map. Let $z = (x, y)$, $\bar{z} = (\bar{x}, \bar{y}) \in S$ be such that $Tz = (T_1 x, T_2 y)$, $T\bar{z} = (T_1 \bar{x}, T_2 \bar{y})$. From hypothesis (H_1) we have

$$\begin{aligned}
 \|T_1 x(t) - T_1 \bar{x}(t)\| &\leq \int_a^t \|f(s, x(s), y(s), \lambda) - f(s, \bar{x}(s), \bar{y}(s), \lambda)\| ds \\
 &\leq \int_a^t p(s) \|z(s) - \bar{z}(s)\|_B \exp(-Ls) \exp(Ls) ds \\
 &\leq \alpha \|z - \bar{z}\|_1 \exp(Lt),
 \end{aligned} \tag{12}$$

and

$$\begin{aligned}
 \|T_2 y(t) - T_2 \bar{y}(t)\| &\leq \int_t^b \|g(s, x(s), y(s), \lambda) - g(s, \bar{x}(s), \bar{y}(s), \lambda)\| ds \\
 &\leq \int_t^b q(s) \|z(s) - \bar{z}(s)\|_B \exp(-Ls) \exp(Ls) ds \\
 &\leq \beta \|z - \bar{z}\|_1 \exp(Lt).
 \end{aligned} \tag{13}$$

From (12) and (13) we have

$$\|Tz - T\bar{z}\|_1 \leq (\alpha + \beta) \|z - \bar{z}\|_1. \tag{14}$$

Since $\alpha + \beta < 1$, it follows from the Banach fixed point theorem that T has a unique fixed point in S . The fixed point of T is however a solution of BVP (1)–(4). This completes the proof of the theorem.

The following theorem determine conditions for continuous dependence of solutions of BVP (1)–(4) on a parameter λ . The idea of the proof is based on the result given by the present author in [10, Theorem 2] (see also [11]).

THEOREM 2. *Let (H_1) , (H_2) and (H_3) hold. Then the solution $(x(t, \lambda), y(t, \lambda))$ of BVP (1)–(4) belonging to S is continuous with respect to the variables (t, λ) in $I \times R$.*

Proof. For $z = (x, y) \in S$ we define the operator $T_\lambda z = (T_{1\lambda} x, T_{2\lambda} y)$, where $T_{1\lambda} x$ and $T_{2\lambda} y$ are defined by the right sides of the equations (8) and (9), respectively. From (14) we have

$$\|T_\lambda z - T_\lambda \bar{z}\|_1 \leq (\alpha + \beta) \|z - \bar{z}\|_1. \tag{15}$$

Next from hypothesis (H_3) we obtain for $\lambda, \lambda_0 \in R$,

$$\begin{aligned}
 \|T_{1\lambda} x(t) - T_{1\lambda_0} x(t)\| &\leq \int_a^t \|f(s, x(s), y(s), \lambda) - f(s, x(s), y(s), \lambda_0)\| ds \\
 &\leq \int_a^t p_0(s) |\lambda - \lambda_0| ds \\
 &\leq M_1 |\lambda - \lambda_0| \exp(Lt),
 \end{aligned} \tag{16}$$

and

$$\begin{aligned} \|T_{2\lambda} y(t) - T_{2\lambda_0} y(t)\| &\leq \int_I^b \|g(s, x(s), y(s), \lambda) - g(s, x(s), y(s), \lambda_0)\| ds \\ &\leq \int_I^b q_0(s) |\lambda - \lambda_0| ds \\ &\leq M_2 |\lambda - \lambda_0| \exp(Lt). \end{aligned} \quad (17)$$

From (16) and (17) we have

$$\|T_\lambda z - T_{\lambda_0} z\|_1 \leq (M_1 + M_2) |\lambda - \lambda_0|. \quad (18)$$

From Theorem 1 there exists a unique function $z(t, \lambda) = (x(t, \lambda), y(t, \lambda))$ such that $T_\lambda z(t, \lambda) = (T_{1\lambda} x(t, \lambda), T_{2\lambda} y(t, \lambda)) = z(t, \lambda)$ and $T_{\lambda_0} z(t, \lambda_0) = (T_{1\lambda_0} x(t, \lambda_0), T_{2\lambda_0} y(t, \lambda_0)) = z(t, \lambda_0)$ for $t \in I$ and $\lambda, \lambda_0 \in R$. From (15) and (18) we have

$$\begin{aligned} \|z(t, \lambda) - z(t, \lambda_0)\|_1 &\leq \|T_\lambda z(t, \lambda) - T_\lambda z(t, \lambda_0)\|_1 \\ &\quad + \|T_\lambda z(t, \lambda_0) - T_{\lambda_0} z(t, \lambda_0)\|_1 \\ &\leq (\alpha + \beta) \|z(t, \lambda) - z(t, \lambda_0)\|_1 + (M_1 + M_2) |\lambda - \lambda_0|, \end{aligned}$$

and hence

$$\|z(t, \lambda) - z(t, \lambda_0)\|_1 \leq (M_1 + M_2) [1 - (\alpha + \beta)]^{-1} |\lambda - \lambda_0|.$$

This shows that the function $z(t, \lambda)$ is continuous with respect to the variable λ in R uniformly with respect to the variable t in I and consequently $z(t, \lambda)$ is also continuous with respect to the two variables (t, λ) in $I \times R$. The proof of the theorem is complete.

We note that BVP (1)–(4) is of more general type and contains as a special case the boundary value problems for second order equations studied by Baily [1]. In [7, 8, 12, 13] the authors have studied BVP (1)–(4) when $\lambda = 0$ by using different techniques. For earlier results on the same type of problems, see [2–5, 14]. An important feature of our approach here is that it is elementary and provides a uniform treatment for problems of this type.

REFERENCES

1. P. B. BAILEY, An eigenvalue theorem for nonlinear second-order differential equations, *J. Math. Anal. Appl.* **20** (1967), 94–102.
2. J. W. BEBERNES AND R. WILHELMSEN, A technique for solving two-dimensional boundary value problems, *SIAM J. Appl. Math.* **17** (1969), 1060–1064.

3. J. W. BEBERNES AND R. WILHELMSEN, A general boundary value problem technique, *J. Differential Equations* **8** (1970), 404–415.
4. S. R. BERNFELD AND V. LAKSHMIKANTHAM, Existence of solutions of ordinary differential equations with generalized boundary conditions, *Trans. Amer. Math. Soc.* **184** (1973), 261–274.
5. S. R. BERNFELD AND V. LAKSHMIKANTHAM, “An Introduction to Nonlinear Boundary Value Problems,” Academic Press, New York, 1974.
6. F. BRAUER, Nonlinear perturbations of Sturm–Liouville boundary value problems, *J. Math. Anal. Appl.* **22** (1968), 591–598.
7. R. E. GAINES AND C. P. GUPTA, Ordinary differential systems of optimal control type with monotone reversed nonlinearities, *J. Math. Anal. Appl.* **64** (1978), 494–504.
8. V. LAKSHMIKANTHAM AND B. G. PACHPATTE, Monotone method for nonlinear boundary value problems arising in transport process, *Appl. Math. Comput.* **9** (1981), 35–41.
9. J. W. MACKI AND P. WALTMAN, A nonlinear boundary value problem of Sturm–Liouville type for a two-dimensional system of ordinary differential equations, *SIAM J. Appl. Math.* **21** (1971), 225–231.
10. B. G. PACHPATTE, On a nonlinear Volterra integral-functional equation, *Funkcial. Ekvac.* **26** (1983), 1–9.
11. B. G. PACHPATTE, On generalized Hammerstein type integral equation, *J. Math. Anal. Appl.* **106** (1985), 85–90.
12. B. G. PACHPATTE, Boundary value problems for nonlinear systems of differential equations in Banach spaces, *Funkcial. Ekvac.* **25** (1982), 199–206.
13. B. G. PACHPATTE AND A. S. VATSALA, Monotone method for equations describing transport phenomena in a Banach space, *Appl. Math. Comput.* **9** (1981), 165–174.
14. P. WALTMAN, Existence and uniqueness of solutions of boundary value problems for two dimensional systems of nonlinear differential equations, *Trans. Amer. Math. Soc.* **153** (1971), 223–234.
15. W. M. WHYBURN, Existence and oscillation theorems for nonlinear differential systems of second order, *Trans. Amer. Math. Soc.* **30** (1928), 848–854.